

Microarrays technique to evaluate the impact of a well-oxygenated brain cells environment or resveratrol on Sirt-1 gene in Wistar rats

GARRIGUE Philippe*, GUILLET Benjamin*, LANDRIER Jean-François**, MOUHAJIR Yassin**,
MERCIER Béatrice***, SEREE Eric**

*CERIMED Marseille France **C2VN INRA INSERM Aix Marseille University Marseille France ***Holiste lab Marcigny, France
Correspondence should be addressed to : bmercier@holiste.com

Introduction



Many circumstances allow the creation of free radicals: aging, pollutions, stress, mitochondrial dysfunction etc. – but also hypoxia (1) and hyperoxia (2).

The aim of this study is to evaluate the **impact of a supplementation in resveratrol (a main antioxidant molecule), or an optimal oxygenation at the brain cellular level** (through regular breathing session of the Bol d'air Jacquier® [BAJ] device) **on gene expression in brain.**

Material and method



Rat's BAJ breathing sessions



Microarrays' technique

The “Bol d'Air Jacquier®” device has been used to allow a well-oxygenated environment at the brain cell level.

The device generates peroxidizing volatile terpenes from organic essential oil of turpentine. It is proved that the peroxidizing molecules are able to optimize the oxygen permeation at the cellular level, during and subsequently to a breathing session (3). So, it is used as an help to fight hypoxia due to environmental pollution, ageing, or inflammatory disease. It is also an help for the control of glycated protein (4) and thus reducing oxidative stress (5).

A major natural molecule, the resveratrol, has been used as a major antioxidant

Resveratrol is a phytochemical with anti-carcinogenic, antioxidant, anti-inflammatory and cardioprotective effects which is found in some dietary sources such as grapes, berries, peanuts or red wine (6). Various galenic form exist to run resveratrol in bodies. We opted for an innovative soluble galenic form able to overcome the usual low absorption of trans-Resveratrol (7).


Microarray technology has been used to assess the genetical impact of a better cell oxygenation or resveratrol treatment on rat's brain.

This technology allows the analysis of the expression of more than 30000 genes, up or down-regulated under the impact of the various treatments (8).

Investigation: on 24 male Wistar rats, 12-month-old and weighing at first between 650 and 750 g, randomised into 3 groups (control, resveratrol and BAJ), fed with resveratrol (50 mg/l) or using BAJ 5 days/week, 5 months.


Results and discussion

Among the up and down-regulated genes, several results sound very interesting and relevant, specially for the Sirt-1 genes.



The BAJ highly up-regulates (around 170%) the Sirt-1 gene expression, but not resveratrol.

Resveratrol doesn't up or down-regulated the genes, but it is proved it increases the activity of the enzymes Sirt-1 - the pathway isn't direct gene activation (9).



Physiological roles of sirtuins (10)

Sirtuin-1 (*silent mating type information regulation 2 homolog*, also known as *NAD-dependent deacetylase sirtuin-1*), is a protein that in humans is encoded by the Sirt-1 gene. These enzymes play a vital role by deacetylate proteins and contribute to cellular regulation (10-13).

Thus, contribute to enhancing the activity of sirtuines is very important for fighting against ageing, especially in brain.

Conclusion

It is the first demonstration of the BAJ effect on the genetic expression in brain. This increase of Sirt-1 expression by BAJ could be impact positively and protect the brain from neurodegenerative diseases. Resveratrol activates the Sirt-1 without modify its gene expression.

The complementary effect of both treatments should be used to fight inflammation and brain ageing.

References

1. Alcaraz-García MJ, Albaladejo MD, Acevedo C, Olea A, Zamora S, Martínez P, Parra S. *Effects of hyperoxia on biomarkers of oxidative stress in closed-circuit oxygen military divers.* J Physiol Biochem. 2008 Jun;64(2):135-41.
2. Miyata T, Takizawa S, van Ypersele de Strihou C. *Hypoxia. 1. Intracellular sensors for oxygen and oxidative stress: novel therapeutic targets.* Am J Physiol Cell Physiol. 2011 Feb;300(2):C226-31.
3. Mercier B. *Bol d'Air Jacquier®: biological and antiradicalizing effects of peroxidizing terpenes* PhD thesis, 2008DIJS019, University of Burgundy 440-16 pp. [in French].
4. Mercier B., Prost J., Prost M. *A well-oxygenated cells environment may help to fight against protein glycation.* IJOMEH 2011;24(1):102-107.
5. Mercier B, Prost J, Prost M. *Antioxidant Activity of Bol d'Air Jacquier® Breathing Sessions in Wistar Rats - First Studies.* Int J Occup Med Environ Health. 2008, Jan 1;21(1):31-46.
6. Ghowsi M, Khazali H, Sisakhtnezhad S. *The effect of resveratrol on oxidative stress in the liver and serum of a rat model of polycystic ovary syndrome: An experimental study.* Int J Reprod Biomed. 2018 Mar;16(3):149-158.
7. Amiot MJ, Romier B, Dao TM, Fanciullino R, Ciccolini J, Burcelin R, Pechere L, Emond C, Savouret JF, Serree E. *Optimization of trans-Resveratrol bioavailability for human therapy.* Biochimie. 2013 Jun;95(6):1233-8. doi: 10.1016/j.biochi.2013.01.008. Epub 2013 Jan 31.
8. Kenichi I, Yoshikuni I, Shuji N *Genomic-Wide Analysis with Microarrays in Human Oncology* Microarrays (Basel). 2015 Dec; 4(4): 454-473.
9. Sun C, Zhang F, Ge X, Yan T, Chen X, Shi X, Zhai Q. *SIRT1 improves insulin sensitivity under insulin-resistant conditions by repressing PTP1B.* Cell Metabolism. 6 (4): 307-19.
10. Finkel T, Deng CX, Mostoslavsky R. *Recent progress in the biology and physiology of sirtuins.* Nature. 2009 Jul 30;460(7255):587-91. doi: 10.1038/nature08197.
11. SIRT1 sirtuin 1 [Homo sapiens (human)] Gene ID: 23411, updated on 27-May-2018
12. Sinclair DA, Guarente L. *Unlocking the Secrets of Longevity Genes* Scientific American. 294 (3): 48-51, 54-7. 2006.
13. Frye RA. *Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity.* Biochemical and Biophysical Research Communications. 260 (1): 273-79, 1999.